



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2017

---

## Host Responses to *Malassezia* spp. in the Mammalian Skin

Sparber, Florian ; LeibundGut-Landmann, Salomé

DOI: <https://doi.org/10.3389/fimmu.2017.01614>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-142333>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Sparber, Florian; LeibundGut-Landmann, Salomé (2017). Host Responses to *Malassezia* spp. in the Mammalian Skin. *Frontiers in Immunology*:8:1614.

DOI: <https://doi.org/10.3389/fimmu.2017.01614>



# Host Responses to *Malassezia* spp. in the Mammalian Skin

Florian Sparber\* and Salomé LeibundGut-Landmann\*

Section of Immunology, Vetsuisse Faculty, University of Zürich, Zürich, Switzerland

## OPEN ACCESS

### Edited by:

Bernhard Hube,  
Hans Knöll Institut, Germany

### Reviewed by:

Ruth Ashbee,  
University of Leeds, United Kingdom  
Annika Elisabet Scheynius,  
Karolinska Institute (KI), Sweden  
Peter Mayer,  
Justus Liebig Universität  
Gießen, Germany

### \*Correspondence:

Florian Sparber  
florian.sparber@uzh.ch;  
Salomé LeibundGut-Landmann  
salome.leibundgut-landmann@uzh.ch

### Specialty section:

This article was submitted to  
Microbial Immunology,  
a section of the journal  
Frontiers in Immunology

**Received:** 28 September 2017

**Accepted:** 08 November 2017

**Published:** 22 November 2017

### Citation:

Sparber F and LeibundGut-Landmann S (2017) Host Responses to *Malassezia* spp. in the Mammalian Skin.  
Front. Immunol. 8:1614.  
doi: 10.3389/fimmu.2017.01614

The skin of mammalian organisms is home for a myriad of microbes. Many of these commensals are thought to have beneficial effects on the host by critically contributing to immune homeostasis. Consequently, dysbiosis can have detrimental effects for the host that may manifest with inflammatory diseases at the barrier tissue. Besides bacteria, fungi make an important contribution to the microbiota and among these, the yeast *Malassezia* widely dominates in most areas of the skin in healthy individuals. There is accumulating evidence that *Malassezia* spp. are involved in a variety of skin disorders in humans ranging from non- or mildly inflammatory conditions such as dandruff and pityriasis versicolor to more severe inflammatory skin diseases like seborrheic eczema and atopic dermatitis. In addition, *Malassezia* is strongly linked to the development of dermatitis and otitis externa in dogs. However, the association of *Malassezia* spp. with such diseases remains poorly characterized. Until now, studies on the fungus–host interaction remain sparse and they are mostly limited to experiments with isolated host cells *in vitro*. They suggest a multifaceted crosstalk of *Malassezia* spp. with the skin by direct activation of the host *via* conserved pattern recognition receptors and indirectly *via* the release of fungus-derived metabolites that can modulate the function of hematopoietic and/or non-hematopoietic cells in the barrier tissue. In this review, we discuss our current understanding of the host response to *Malassezia* spp. in the mammalian skin.

**Keywords:** *Malassezia*, commensalism, opportunistic pathogenic fungi, skin disorders, innate immunity, adaptive immunity, allergic response, indoles

## INTRODUCTION

*Malassezia* spp. are lipophilic yeasts, which are part of the skin microbiota of many mammals and birds. In fact, the genus *Malassezia* is by far the most abundant eukaryotic member of the microbial flora of the skin in these organisms (1). Most *Malassezia* spp. have a predilection for seborrheic skin sites such as the scalp and the trunk. They rely on exogenous fatty acid sources for their nutritive requirements because of their lack of genes encoding for the fatty acid synthase and genes involved in carbohydrate metabolism (2–4). In agreement, the cell wall of *Malassezia* spp. is particularly rich in lipids (5).

The genus *Malassezia* currently comprises 17 species, three of which have only recently been proposed (6–8). *Malassezia globosa*, *Malassezia restricta*, and *Malassezia sympodialis* are most frequently isolated from the healthy human skin with distinct relative frequencies at specific body sites (1, 9). The age of the host and geographic factors also influence their distribution (10). *Malassezia pachydermatis*, *Malassezia nana*, and *Malassezia caprae* are found predominantly in non-human hosts (6). Surprisingly, the microbial communities of the skin are astonishingly stable and

maintained over time, despite the skin's exposure to the external environment (11). It is currently unknown whether *Malassezia* spp. play a mutualistic role and may thus contribute to immune homeostasis of the host.

Apart from their commensal nature, *Malassezia* spp. are also associated with common skin disorders such as pityriasis versicolor and seborrheic dermatitis as well as more severe inflammatory skin pathologies including atopic eczema and atopic dermatitis in humans (10) and dermatitis and otitis externa in animals, most frequently in dogs (12). The composition of the skin mycobiome can vary under pathological conditions and some species of *Malassezia* such as *M. sympodialis* and *Malassezia furfur* are found to be enriched in certain skin disorders (10). To date, a causative link between *Malassezia* and disease development has only been made for Pityriasis versicolor, while the role of the yeast in other pathologies remains correlative (10, 13, 14). Changes in the degree of colonization in diseased compared to healthy skin have been documented in dogs (15) but remain uncertain in humans (16).

The pathophysiology of *Malassezia*-associated skin conditions is largely unknown. The lack of knowledge on the cellular and molecular interactions between *Malassezia* spp. and the host preclude a better understanding of the factors determining commensalism versus disease. Herein, we review the current knowledge with regard to how the host recognizes *Malassezia* spp. and responds to it (Figure 1).


## SENSING OF *Malassezia* spp. BY THE HOST

Through their localization in the skin, *Malassezia* spp. interact primarily with keratinocytes, tissue-resident dendritic cells (DCs), and macrophages, as well as with myeloid cells that are recruited to the skin under inflammatory conditions. Activation of DCs is key for induction of adaptive immunity and memory formation. The fungus is recognized by the host either directly through interaction of fungal cell wall components with membrane bound pattern recognition receptors (PRRs) or indirectly through soluble metabolites that are released by *Malassezia* spp. The set of receptors expressed by the hematopoietic and the non-hematopoietic compartment are largely distinct.

### Direct Recognition of *Malassezia* spp. by Surface-Bound Receptors

The fungal cell wall is rich in carbohydrates and glycoproteins that are recognized by PRRs of the family of Syk-coupled C-type lectin receptor (CLR), which are expressed primarily by myeloid cells (17, 18). Binding to these receptors results in ligand internalization and activation of multiple signaling pathways, including the MAPK, NF- $\kappa$ B, and NFAT pathways as well as the inflammasome.

The polysaccharides of the *Malassezia* cell wall are organized differently than in other fungal species analyzed to date (19, 20).



**FIGURE 1 |** Interaction of *Malassezia* spp. with the mammalian skin. Direct interactions involve various PRRs, which recognize fungal cell wall constituents and are distinctly expressed on the surface of non-hematopoietic (i.e., keratinocytes) and hematopoietic cells (i.e., Langerhans cells, mast cells) of the skin. Spatial and temporal signal integration of different PRR signals results in the induction of inflammation and immunity or, alternatively, in the regulation and tolerance of the host toward *Malassezia* spp. Indirect interactions of *Malassezia* spp. with the skin include fungus-derived metabolites such as irritant fatty acids on the one hand and indoles that are potent agonists for the AhR, which is expressed by various skin cells, on the other hand. KCs, keratinocytes; LCs, Langerhans cells; DCs, dendritic cells; Mφ, macrophages; MCs, mast cells; SC, stratum corneum; AhR, aryl hydrocarbon receptor.

Moreover, the cell wall is surrounded by a lipid-rich outer layer (21). Several CLRs have been shown to respond to *Malassezia* spp. *in vitro*. The two FcRγ-associated receptors Dectin-2 and Mincle sense *Malassezia* spp., albeit through recognition of distinct ligands (22). While Mincle binds to two distinct glycolipids in *Malassezia*, Dectin-2 recognizes the fungus through α-1,2-linked mannose. High-mannose binding is a general feature of Dectin-2, which is reported to recognize a variety of fungi, including *Candida albicans*, *Saccharomyces cerevisiae*, *Blastomyces dermatitidis*, *Aspergillus fumigatus*, *Cryptococcus neoformans*, and *Fonsecaea pedrosoi* (23). In contrast, *Malassezia* spp. were initially found to be unique agonists of Mincle when a large panel of 50 different fungi was tested in a glycoconjugate microarray (24). More recently, other fungi such as *Pneumocystis carinii*, *F. pedrosoi*, and *Fonsecaea monomorpha* were also reported to engage Mincle (25–27), in addition to bacterial ligands (28–32), mammalian alarmins released from damaged cells (33, 34) and even cholesterol crystals (35, 36). Mincle is thus a highly pleiotropic receptor, which can bind chemically and structurally distinct ligands through at least two complementary binding sites (37–40). The β-glucan receptor Dectin-1, which was the first member of the family of Syk-coupled CLRs to be identified (41), was also found to sense *Malassezia* and was linked to the activation of the NLRP3 inflammasome (42). Finally, Langerin was suggested to act as a receptor for *Malassezia* in the skin due to its prominent expression by epidermal Langerhans cells and by a subset of dermal DCs. Direct binding of the fungus to recombinant Langerin was indeed observed (43, 44).

Activation of myeloid cells by *Malassezia* spp. via these different CLRs was shown to induce the secretion of proinflammatory cytokines. However, the relative contribution of individual receptors to fungal control *in vivo* during commensalism and in infectious settings remains to be determined. At least partial redundancy of receptors that signal via the same pathway may occur, similarly to what was found for other fungi (45, 46). Dissecting the role of Mincle in the context of *Malassezia* spp. in more detail will also be interesting in light of its reported antagonizing activity, e.g., in response to *Fonsecaea* spp. (27), and thus this receptor may also mediate regulatory or inhibitory responses to *Malassezia* spp.

In addition to CLRs, Toll-like receptors (TLRs), and in particular TLR2, also contribute to fungal recognition by the host. TLR2 was implicated in sensing of *Malassezia* spp. and inducing a proinflammatory response characterized by the release of cytokines, chemokines and antimicrobial peptides by keratinocytes (47–50).

The proinflammatory response is generally enhanced by lipid removal from the yeast to enhance exposure of fungal cell wall carbohydrates (51, 52). In contrast, thymic stromal lymphopoietin secretion from keratinocytes was found to be induced specifically by the lipid layer components of *M. restricta* and *M. globosa* but not by yeasts that were depleted of lipids (53).

## Indirect Interaction

Specific products of *Malassezia* metabolic pathways are thought to act as virulence factors promoting inflammation and pathology, while others downregulate the production of inflammatory mediators and thereby contribute to immune regulation. Fungal

strains with altered production of such factors have been linked to *Malassezia*-associated skin disorders (54–56).

*Malassezia*-derived lipases and phospholipases, which are required to assimilate host-derived lipids, can initiate an inflammatory response in the skin by releasing unsaturated free fatty acids from the sebum lipids (57–60). Oleic acid has irritant and desquamative effects on keratinocytes (61–63), whereas arachidonic acid produces proinflammatory eicosanoids and leads to inflammation and damage to the stratum corneum, thereby contributing to the disruption of the epithelial barrier function and induction of abnormal keratinization (64).

*Malassezia furfur* is able to convert tryptophan into a variety of indole alkaloids. This pathway is mainly active if tryptophan is the sole source of nitrogen (65). *M. furfur*-derived indoles including malassezin, indirubin, and indolo [3,2-b] carbazole (ICZ) serve as potent ligands for the host aryl hydrocarbon receptor (AhR) and thereby potentially modify the function of all cells in the epidermis expressing this receptor (54, 55, 66, 67). For example, some tryptophan metabolites can promote apoptosis of melanocytes (68) or inhibit the respiratory burst in neutrophils (69). Given the broad spectrum of biological responses that are influenced by AhR activity, *M. furfur* may engage this pathway to modulate inflammation and/or promote skin immune homeostasis (70) but may also promote skin pathology (71) or even contribute to carcinogenesis (72). The significance of yeast-derived indoles in each of these contexts remains to be demonstrated *in vivo*.

## INNATE IMMUNITY TO *Malassezia* spp.

The majority of what is currently known about the host response to *Malassezia* spp. is based on *in vitro* studies with isolated myeloid cells or keratinocyte cell lines. Stimulation of these cells with *Malassezia* yeast leads to the induction of mainly proinflammatory cytokines, chemokines, and antimicrobial peptides (22, 24, 47–52, 73–76). In line with an inflammatory character of the innate response to the fungus, the intraperitoneal injection of *Malassezia* into mice results in the recruitment of neutrophils to the peritoneum (24). Only few studies have examined regulatory cytokines such as IL-10 and TGF-β by the yeast (24, 49, 51, 74, 77), but these may be relevant with regard to the role of *Malassezia* spp. as a skin commensal.

Given the association of *Malassezia* spp. with inflammatory skin disorders and allergic responses, the fungus may also interact with mast cells. Progenitor cell-derived mast cells from atopic patients show increased release of proinflammatory cytokines upon stimulation with *Malassezia* (76) and are enriched in the skin of atopic eczema patients where they are positioned in the superficial dermis and can interact with the fungus (78). Mast cell activation in response to *Malassezia* spp. has also been reported in studies with bone-marrow-derived mast cells. These cells are directly activated by the fungus in a TLR2-dependent manner and release inflammatory mediators and cytokines (79). Moreover, the crosslinking of the high-affinity IgE receptor (FcεRI) by antigen-bound IgE can induce mast cell degranulation (79). Therefore, mast cells may contribute to further barrier disruption and thereby amplify the inflammatory response.



The access of *Malassezia* to immune cells in the skin may be facilitated by disruption of the epithelial barrier as it frequently occurs during chronic inflammation. Moreover, *Malassezia* spp. were reported to release nanovesicles/exosomes that contain immunogenic proteins and trigger increased release of cytokines by DCs (80).

## ADAPTIVE IMMUNITY TO *Malassezia* spp.

As a commensal, *Malassezia* interacts continuously with the immune system. Therefore, cellular and humoral immune memory to the fungus can be evidenced in healthy individuals (81). Although there are fewer studies related to dogs when compared with humans, dogs also develop cellular and humoral immune responses to their commensal yeast, *M. pachydermatis* (82–84). Generally, the adaptive immune responses are heightened and qualitatively distinct in patients with *Malassezia*-associated diseases.

### Humoral Responses

During steady state, *Malassezia*-specific antibodies are predominantly of the IgG and IgM isotypes (81). In contrast, although *Malassezia*-specific IgE is not usually detected in healthy individuals, it is common in atopic patients (85). A positive correlation was found between the sensitization to *Malassezia*-specific IgE and the severity of atopic dermatitis (86, 87). Similar observations were made in atopic dogs (83, 84). However, whether the IgE response plays a pathogenic role in atopic and other *Malassezia*-associated inflammatory disorders or rather serves as a marker for the severity of disease remains unclear.

### T Cell Responses

Patients with atopic dermatitis often show positive skin prick test and atopic patch test reactions to *Malassezia* (85). T cell-responsiveness to *Malassezia* in such patients was associated with a Th2 response (88), in line with the classical paradigm of Th2-polarized allergic T cells. GATA3<sup>+</sup> T cells were identified in pityriasis versicolor lesions (89) and likewise *Malassezia*-specific T cell in allergic dogs were found to be strongly polarized toward a type 2 response (82). More recently, other T helper cell subsets such as Th17 and Th22 cells have been found enriched in allergic individuals (90, 91) as well as in non-allergic immune-mediated skin diseases such as psoriasis (92). Consistent with this notion, *Malassezia*-reactive skin homing T cells from *Malassezia*-sensitized atopic dermatitis patients comprise not only Th1 and Th2 subsets but also IL-17- and IL-22-secreting cells (93). Of note, IL-4/IL-17 coproducers have also been described in the context of atopic eczema especially in children (94). Importantly, Th17 differentiation is a hallmark of T cell responses induced by CLR signaling (46) and T cells directed against other fungi, in particular *Candida* spp., belong predominantly to the Th17 subset (95). Whether and how IL-17 and/or IL-22 may contribute to pathogenicity in atopic dermatitis remains to be determined. It is also unknown to which subset *Malassezia*-specific T cells belong in healthy individuals and to what extent T cell plasticity contributes to sensitization.

## *Malassezia* Allergens

To date, 13 *Malassezia*-derived allergens have been identified from *M. furfur* and *M. sympodialis* (3, 96). Interestingly, more allergens are released from *M. sympodialis* when cultured at the increased pH conditions of atopic skin compared with culture at the pH of healthy skin (97). Several of the known allergens belong to a class of phylogenetically highly conserved proteins and display a high degree of homology with the corresponding mammalian proteins. Cross-reactivity between *Malassezia*-derived allergens and endogenous human proteins (e.g., thioredoxin, manganese-dependent superoxide dismutase) has been indeed demonstrated (93, 98, 99). Therefore, the induction of autoreactive T cells by *Malassezia* allergens may play a role in sustained inflammation.

## CONCLUSION

*Malassezia* spp. have been implicated in various pathologies. Yet, direct evidence for a causal relationship between *Malassezia* spp. and the mammalian host remains elusive. For instance, it is unclear whether *Malassezia* actively promotes atopic dermatitis or whether the inflammatory environment in the atopic skin triggers a dysregulated immune response toward the fungus.

At the basis of this is the key question of what determines the balance between commensalism and pathogenicity of *Malassezia* spp. The answer likely relates to changes occurring in both the fungus (55) (e.g., variable secretion of AhR agonists) and in the host (e.g., barrier defects, changes in immune polarization) which are responsible for promoting the development of pathology. Changes in the environment such as seasonal variations in sebum production have also been linked to altered disease prevalence (100).

Inter-species variations in the skin mycobiome may further contribute as different species of *Malassezia* can induce variable inflammatory responses (51, 75, 101). Moreover, *Malassezia* spp. have been shown to display a large intra-species diversity (73) similarly to what is known for other opportunistic fungal pathogens (102), and thus the exact composition of *Malassezia* strains and species present in an individual at a given time may contribute to different outcomes in the interaction between the fungus and the host. The recently completed assembly and detailed annotation of the genome of *M. sympodialis* makes an important contribution to approach this complexity (103). Future research will help fill the important gaps in our knowledge on the pathophysiology of and the host response to *Malassezia* *in vivo*. Enhanced understanding of host-*Malassezia* interactions may contribute to improved diagnostic and therapeutic options for patients affected by *Malassezia*-associated pathologies.

## AUTHOR CONTRIBUTIONS

Both authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

## FUNDING

This work was supported by the University of Zürich.

## REFERENCES

- Findley K, Oh J, Yang J, Conlan S, Deming C, Meyer JA, et al. Topographic diversity of fungal and bacterial communities in human skin. *Nature* (2013) 498:367–70. doi:10.1038/nature12171
- Wu G, Zhao H, Li C, Rajapakse MP, Wong WC, Xu J, et al. Genus-wide comparative genomics of *Malassezia* delineates its phylogeny, physiology, and niche adaptation on human skin. *PLoS Genet* (2015) 11:e1005614. doi:10.1371/journal.pgen.1005614
- Gioti A, Nystedt B, Li W, Xu J, Andersson A, Averette AF, et al. Genomic insights into the atopic eczema-associated skin commensal yeast *Malassezia sympodialis*. *mBio* (2013) 4:e00572–12. doi:10.1128/mBio.00572-12
- Park M, Cho Y-J, Lee YW, Jung WH. Whole genome sequencing analysis of the cutaneous pathogenic yeast *Malassezia restricta* and identification of the major lipase expressed on the scalp of patients with dandruff. *Mycoses* (2017) 60:188–97. doi:10.1111/myc.12586
- Thompson E, Colvin JR. Composition of the cell wall of *Pityrosporum ovale* (Bizzozero) Castellani and Chalmers. *Can J Microbiol* (1970) 16:263–5. doi:10.1139/m70-048
- Cabanes FJ. *Malassezia* yeasts: how many species infect humans and animals? *PLoS Pathog* (2014) 10:e1003892. doi:10.1371/journal.ppat.1003892
- Honnarav P, Prasad GS, Ghosh A, Dogra S, Handa S, Rudramurthy SM. *Malassezia arunalokei* sp. nov., a novel yeast species isolated from seborrheic dermatitis patients and healthy individuals from India. *J Clin Microbiol* (2016) 54:1826–34. doi:10.1128/JCM.00683-16
- Cabanes FJ, Coutinho SDA, Puig L, Bragulat MR, Castella G. New lipid-dependent *Malassezia* species from parrots. *Rev Iberoam Micol* (2016) 33:92–9. doi:10.1016/j.riam.2016.03.003
- Jo J-H, Kennedy EA, Kong HH. Topographical and physiological differences of the skin mycobiome in health and disease. *Virulence* (2017) 8:324–33. doi:10.1080/21505594.2016.1249093
- Prohic A, Jovovic Sadikovic T, Krupalija-Fazlic M, Kuskunovic-Vlahovljak S. *Malassezia* species in healthy skin and in dermatological conditions. *Int J Dermatol* (2016) 55:494–504. doi:10.1111/ijd.13116
- Oh J, Byrd AL, Park M, Kong HH, Segre JA. Temporal stability of the human skin microbiome. *Cell* (2016) 165:854–66. doi:10.1016/j.cell.2016.04.008
- Bond R, Guillot J, Cabanes FJ. *Malassezia* yeasts in animal disease. In: Boekhout T, Mayser P, Guého-Kellermann E, Velegraki A, editors. *Malassezia and the Skin: Science and Clinical Practice*. Berlin, Heidelberg: Springer (2010). p. 271–99.
- Johansson C, Eshaghi H, Linder MT, Jakobson E, Scheynius A. Positive atopy patch test reaction to *Malassezia furfur* in atopic dermatitis correlates with a T helper 2-like peripheral blood mononuclear cells response. *J Invest Dermatol* (2002) 118:1044–51. doi:10.1046/j.1523-1747.2002.01758.x
- Casagrande BF, Fluckiger S, Linder MT, Johansson C, Scheynius A, Cramer R, et al. Sensitization to the yeast *Malassezia sympodialis* is specific for extrinsic and intrinsic atopic eczema. *J Invest Dermatol* (2006) 126:2414–21. doi:10.1038/sj.jid.5700431
- Cafarchia C, Gallo S, Romito D, Capelli G, Chermette R, Guillot J, et al. Frequency, body distribution, and population size of *Malassezia* species in healthy dogs and in dogs with localized cutaneous lesions. *J Vet Diagn Invest* (2005) 17:316–22. doi:10.1177/104063870501700403
- Sandstrom Falk MH, Tengvall Linder M, Johansson C, Bartosik J, Back O, Sarnhult T, et al. The prevalence of *Malassezia* yeasts in patients with atopic dermatitis, seborrheic dermatitis and healthy controls. *Acta Derm Venereol* (2005) 85:17–23. doi:10.1080/00015550410022276
- Underhill DM, Pearlman E. Immune interactions with pathogenic and commensal fungi: a two-way street. *Immunity* (2015) 43:845–58. doi:10.1016/j.immuni.2015.10.023
- Dambua IM, Brown GD. C-type lectins in immunity: recent developments. *Curr Opin Immunol* (2015) 32:21–7. doi:10.1016/j.coi.2014.12.002
- Stalhberger T, Simenel C, Clavaud C, Eijssink VGH, Jourdain R, Delepierre M, et al. Chemical organization of the cell wall polysaccharide core of *Malassezia restricta*. *J Biol Chem* (2014) 289:12647–12656.
- Kruppa MD, Lowman DW, Chen YH, Selander C, Scheynius A, Monteiro MA, et al. Identification of (1 → 6)-β-D-glucan as the major carbohydrate component of the *Malassezia sympodialis* cell wall. *Carbohydr Res* (2009) 344:2474–9. doi:10.1016/j.carres.2009.09.029
- Mittag H. Fine structural investigation of *Malassezia furfur*. II. The envelope of the yeast cells. *Mycoses* (1995) 38:13–21. doi:10.1111/j.1439-0507.1995.tb00003.x
- Ishikawa T, Itoh F, Yoshida S, Saijo S, Matsuzawa T, Gonoi T, et al. Identification of distinct ligands for the C-type lectin receptors Mincle and Dectin-2 in the pathogenic fungus *Malassezia*. *Cell Host Microbe* (2013) 13:477–88. doi:10.1016/j.chom.2013.03.008
- Brown GD, Crocker PR. Lectin receptors expressed on myeloid cells. *Microbiol Spectr* (2016) 4(5):MCHD-0036-2016. doi:10.1128/microbiolspec.MCHD-0036-2016
- Yamasaki S, Matsumoto M, Takeuchi O, Matsuzawa T, Ishikawa E, Sakuma M, et al. C-type lectin Mincle is an activating receptor for pathogenic fungus, *Malassezia*. *Proc Natl Acad Sci U S A* (2009) 106:1897–902. doi:10.1073/pnas.0805177106
- Kottom TJ, Hebrink DM, Jensen PE, Nandakumar V, Wuthrich M, Wang H, et al. The interaction of pneumocystis with the C-type lectin receptor Mincle exerts a significant role in host defense against infection. *J Immunol* (2017) 198:3515–25. doi:10.4049/jimmunol.1600744
- Sousa MdG, Reid DM, Schweighoffer E, Tybulewicz V, Ruland J, Langhorne J, et al. Restoration of pattern recognition receptor costimulation to treat chromoblastomycosis, a chronic fungal infection of the skin. *Cell Host Microbe* (2011) 9:436–43. doi:10.1016/j.chom.2011.04.005
- Wevers BA, Kaptein TM, Zijlstra-Willems EM, Theelen B, Boekhout T, Geijtenbeek TBH, et al. Fungal engagement of the C-type lectin mincle suppresses dectin-1-induced antifungal immunity. *Cell Host Microbe* (2014) 15:494–505. doi:10.1016/j.chom.2014.03.008
- Iborra S, Martinez-Lopez M, Cueto FJ, Conde-Garrosa R, Del Fresno C, Izquierdo HM, et al. *Leishmania* uses Mincle to target an inhibitory ITAM signaling pathway in dendritic cells that dampens adaptive immunity to infection. *Immunity* (2016) 45:788–801. doi:10.1016/j.immuni.2016.09.012
- Rabes A, Zimmermann S, Reppe K, Lang R, Seeberger PH, Suttrop N, et al. The C-type lectin receptor Mincle binds to *Streptococcus pneumoniae* but plays a limited role in the anti-pneumococcal innate immune response. *PLoS One* (2015) 10:e0117022. doi:10.1371/journal.pone.0117022
- Devi S, Rajakumara E, Ahmed N. Induction of Mincle by *Helicobacter pylori* and consequent anti-inflammatory signaling denote a bacterial survival strategy. *Sci Rep* (2015) 5:15049. doi:10.1038/srep15049
- Schick J, Etschel P, Bailo R, Ott L, Bhatt A, Lepenies B, et al. Toll-like receptor 2 and Mincle cooperatively sense corynebacterial cell wall glycolipids. *Infect Immun* (2017) 85:e00075–17. doi:10.1128/IAI.00075-17
- Matsunaga I, Moody DB. Mincle is a long sought receptor for mycobacterial cord factor. *J Exp Med* (2009) 206:2865–8. doi:10.1084/jem.20092533
- Yamasaki S, Ishikawa E, Sakuma M, Hara H, Ogata K, Saito T. Mincle is an ITAM-coupled activating receptor that senses damaged cells. *Nat Immunol* (2008) 9:1179–88. doi:10.1038/ni.1651
- Nagata M, Izumi Y, Ishikawa E, Kiyotake R, Doi R, Iwai S, et al. Intracellular metabolite beta-glucosylceramide is an endogenous Mincle ligand possessing immunostimulatory activity. *Proc Natl Acad Sci U S A* (2017) 114:E3285–94. doi:10.1073/pnas.1618133114
- Kiyotake R, Oh-Hora M, Ishikawa E, Miyamoto T, Ishibashi T, Yamasaki S. Human Mincle binds to cholesterol crystals and triggers innate immune responses. *J Biol Chem* (2015) 290:25322–32. doi:10.1074/jbc.M115.645234
- Kostarnoy AV, Gancheva PG, Lepenies B, Tukhvatulin AI, Dzharullaeva AS, Polyakov NB, et al. Receptor Mincle promotes skin allergies and is capable of recognizing cholesterol sulfate. *Proc Natl Acad Sci U S A* (2017) 114:E2758–65. doi:10.1073/pnas.1611665114
- Feinberg H, Jegouzo SAF, Rowntree TJW, Guan Y, Brash MA, Taylor ME, et al. Mechanism for recognition of an unusual mycobacterial glycolipid by the macrophage receptor mincle. *J Biol Chem* (2013) 288:28457–65. doi:10.1074/jbc.M113.497149
- Jegouzo SAF, Harding EC, Acton O, Rex MJ, Fadden AJ, Taylor ME, et al. Defining the conformation of human mincle that interacts with mycobacterial trehalose dimycolate. *Glycobiology* (2014) 24:291–300. doi:10.1093/glycob/cwu072
- Feinberg H, Rambaruth NDS, Jegouzo SAF, Jacobsen KM, Djurhuus R, Poulsen TB, et al. Binding sites for acylated trehalose analogs of glycolipid

- ligands on an extended carbohydrate recognition domain of the macrophage receptor Mincle. *J Biol Chem* (2016) 291:21222–33.
40. Furukawa A, Kamishikiryō J, Mori D, Toyonaga K, Okabe Y, Toji A, et al. Structural analysis for glycolipid recognition by the C-type lectins Mincle and MCL. *Proc Natl Acad Sci U S A* (2013) 110:17438–43. doi:10.1073/pnas.1312649110
  41. Brown GD, Gordon S. Immune recognition. A new receptor for beta-glucans. *Nature* (2001) 413:36–7. doi:10.1038/35092620
  42. Kistowska M, Fenini G, Jankovic D, Feldmeyer L, Kerl K, Bosshard P, et al. *Malassezia* yeasts activate the NLRP3 inflammasome in antigen-presenting cells via Syk-kinase signalling. *Exp Dermatol* (2014) 23:884–9. doi:10.1111/exd.12552
  43. de Jong MA, Vriend LE, Theelen B, Taylor ME, Fluitsma D, Boekhout T, et al. C-type lectin Langerin is a beta-glucan receptor on human Langerhans cells that recognizes opportunistic and pathogenic fungi. *Mol Immunol* (2010) 47:1216–25. doi:10.1016/j.molimm.2009.12.016
  44. Tateno H, Ohnishi K, Yabe R, Hayatsu N, Sato T, Takeya M, et al. Dual specificity of Langerin to sulfated and mannoseylated glycans via a single C-type carbohydrate recognition domain. *J Biol Chem* (2010) 285:6390–400. doi:10.1074/jbc.M109.041863
  45. Robinson MJ, Osorio F, Rosas M, Freitas RP, Schweighoffer E, Gross O, et al. Dectin-2 is a Syk-coupled pattern recognition receptor crucial for Th17 responses to fungal infection. *J Exp Med* (2009) 206:2037–51. doi:10.1084/jem.20082818
  46. LeibundGut-Landmann S, Gross O, Robinson MJ, Osorio F, Slack EC, Tsoni SV, et al. Syk- and CARD9-dependent coupling of innate immunity to the induction of T helper cells that produce interleukin 17. *Nat Immunol* (2007) 8:630–8. doi:10.1038/ni1460
  47. Angrisano T, Pero R, Paoletti I, Keller S, Lembo L, Baroni A, et al. Epigenetic regulation of IL-8 and beta-defensin genes in human keratinocytes in response to *Malassezia furfur*. *J Invest Dermatol* (2013) 133:2104–2104. doi:10.1038/jid.2013.143
  48. Buommino E, De Filippis A, Parisi A, Nizza S, Martano M, Iovane G, et al. Innate immune response in human keratinocytes infected by a feline isolate of *Malassezia pachydermatis*. *Vet Microbiol* (2013) 163:90–6. doi:10.1016/j.vetmic.2012.12.001
  49. Donnarumma G, Perfetto B, Paoletti I, Oliviero G, Clavard C, Del Bufalo A, et al. Analysis of the response of human keratinocytes to *Malassezia globosa* and *restricta* strains. *Arch Dermatol Res* (2014) 306:763–8. doi:10.1007/s00403-014-1479-1
  50. Baroni A, Orlando M, Donnarumma G, Farro P, Iovene MR, Tufano MA, et al. Toll-like receptor 2 (TLR2) mediates intracellular signalling in human keratinocytes in response to *Malassezia furfur*. *Arch Dermatol Res* (2006) 297:280–8. doi:10.1007/s00403-005-0594-4
  51. Thomas DS, Ingham E, Bojar RA, Holland KT. In vitro modulation of human keratinocyte pro- and anti-inflammatory cytokine production by the capsule of *Malassezia* species. *FEMS Immunol Med Microbiol* (2008) 54:203–14. doi:10.1111/j.1574-695X.2008.00468.x
  52. Kesavan S, Holland KT, Ingham E. The effects of lipid extraction on the immunomodulatory activity of *Malassezia* species in vitro. *Med Mycol* (2000) 38:239–47. doi:10.1080/mmy.38.3.239.247
  53. Ishibashi Y, Sugawara K, Sugita T, Nishikawa A. Secretion of thymic stromal lymphopoietin from human keratinocytes in response to *Malassezia* yeasts. *J Dermatol Sci* (2011) 62:124–8. doi:10.1016/j.jdermsci.2011.02.012
  54. Magiatis P, Pappas P, Gaitanis G, Mexia N, Melliou E, Galanou M, et al. *Malassezia* yeasts produce a collection of exceptionally potent activators of the Ah (dioxin) receptor detected in diseased human skin. *J Invest Dermatol* (2013) 133:2023–30. doi:10.1038/jid.2013.92
  55. Gaitanis G, Magiatis P, Stathopoulou K, Bassukas ID, Alexopoulos EC, Velegraki A, et al. AhR ligands, malassezin, and indolo[3,2-b]carbazole are selectively produced by *Malassezia furfur* strains isolated from seborrheic dermatitis. *J Invest Dermatol* (2008) 128:1620–5. doi:10.1038/sj.jid.5701252
  56. Cafarchia C, Otranto D. Association between phospholipase production by *Malassezia pachydermatis* and skin lesions. *J Clin Microbiol* (2004) 42:4868–9. doi:10.1128/JCM.42.10.4868-4869.2004
  57. Riciputo RM, Oliveri S, Micali G, Sapuppo A. Phospholipase activity in *Malassezia furfur* pathogenic strains. *Mycoses* (1996) 39:233–5. doi:10.1111/j.1439-0507.1996.tb00131.x
  58. Plotkin LI, Squiquera L, Mathov I, Galimberti R, Leoni J. Characterization of the lipase activity of *Malassezia furfur*. *J Med Vet Mycol* (1996) 34:43–8. doi:10.1080/02681219680000071
  59. Lee YW, Lee SY, Lee Y, Jung WH. Evaluation of expression of lipases and phospholipases of *Malassezia restricta* in patients with seborrheic dermatitis. *Ann Dermatol* (2013) 25:310–4. doi:10.5021/ad.2013.25.3.310
  60. Ortiz G, Martin MC, Carrillo-Muñoz AJ, Paya MJ. [Phospholipase and proteinase production by *Malassezia pachydermatis* isolated in dogs with and without otitis]. *Rev Iberoam Micol* (2013) 30:235–8. doi:10.1016/j.riam.2013.01.006
  61. DeAngelis YM, Gemmer CM, Kaczvinsky JR, Kenneally DC, Schwartz JR, Dawson TL Jr. Three etiologic facets of dandruff and seborrheic dermatitis: *Malassezia* fungi, sebaceous lipids, and individual sensitivity. *J Invest Dermatol Symp* (2005) 10:295–7. doi:10.1111/j.1087-0024.2005.10119.x
  62. Katsuta Y, Iida T, Hasegawa K, Inomata S, Denda M. Function of oleic acid on epidermal barrier and calcium influx into keratinocytes is associated with N-methyl D-aspartate-type glutamate receptors. *Br J Dermatol* (2009) 160:69–74. doi:10.1111/j.1365-2133.2008.08860.x
  63. Katsuta Y, Iida S, Fau-Inomata T, Inomata M, Fau-Denda S, Denda M. Unsaturated fatty acids induce calcium influx into keratinocytes and cause abnormal differentiation of epidermis. *J Invest Dermatol* (2005) 124:1008–13. doi:10.1111/j.0022-202X.2005.23682.x
  64. Plotkin LI, Mathov I, Squiquera L, Leoni J. Arachidonic acid released from epithelial cells by *Malassezia furfur* phospholipase A2: a potential pathophysiological mechanism. *Mycologia* (1998) 90:163–9. doi:10.2307/3761291
  65. Mayser P, Wille G, Imkamp A, Thoma W, Arnold N, Monsees T. Synthesis of fluorochromes and pigments in *Malassezia furfur* by use of tryptophan as the single nitrogen source. *Mycoses* (1998) 41:265–71. doi:10.1111/j.1439-0507.1998.tb00336.x
  66. Mexia N, Gaitanis G, Velegraki A, Soshilov A, Denison MS, Magiatis P. Pityriazepin and other potent AhR ligands isolated from *Malassezia furfur* yeast. *Arch Biochem Biophys* (2015) 571:16–20. doi:10.1016/j.abb.2015.02.023
  67. Mayer P, Schafer U, Kramer HJ, Irlinger B, Steglich W. Pityriacitrin – an ultraviolet-absorbing indole alkaloid from the yeast *Malassezia furfur*. *Arch Dermatol Res* (2002) 294:131–4. doi:10.1007/s00403-002-0294-2
  68. Kramer HJ, Podobinska M, Bartsch A, Battmann A, Thoma W, Bernd A, et al. Malassezin, a novel agonist of the aryl hydrocarbon receptor from the yeast *Malassezia furfur*, induces apoptosis in primary human melanocytes. *Chembiochem* (2005) 6:860–5. doi:10.1002/cbic.200400247
  69. Kramer HJ, Kessler D, Hipler UC, Irlinger B, Hort W, Bodeker RH, et al. Pityriarubins, novel highly selective inhibitors of respiratory burst from cultures of the yeast *Malassezia furfur*: comparison with the bisindolylmaleimide arcyriarubin A. *Chembiochem* (2005) 6:2290–7. doi:10.1002/cbic.200500163
  70. Vlachos C, Schulte BM, Magiatis P, Adema GJ, Gaitanis G. *Malassezia*-derived indoles activate the aryl hydrocarbon receptor and inhibit toll-like receptor-induced maturation in monocyte-derived dendritic cells. *Br J Dermatol* (2012) 167:496–505. doi:10.1111/j.1365-2133.2012.11014.x
  71. Tsuchi M, Hida A, Negishi T, Katsuoka F, Noda S, Mimura J, et al. Constitutive expression of aryl hydrocarbon receptor in keratinocytes causes inflammatory skin lesions. *Mol Cell Biol* (2005) 25:9360–8. doi:10.1128/MCB.25.21.9360-9368.2005
  72. Gaitanis G, Velegraki A, Magiatis P, Pappas P, Bassukas ID. Could *Malassezia* yeasts be implicated in skin carcinogenesis through the production of aryl-hydrocarbon receptor ligands? *Med Hypotheses* (2011) 77:47–51. doi:10.1016/j.mehy.2011.03.020
  73. Buommino E, Nocera FP, Parisi A, Rizzo A, Donnarumma G, Mallardo K, et al. Correlation between genetic variability and virulence factors in clinical strains of *Malassezia pachydermatis* of animal origin. *New Microbiol* (2016) 39:216–23.
  74. Donnarumma G, Paoletti I, Buommino E, Orlando M, Tufano MA, Baroni A. *Malassezia furfur* induces the expression of beta-defensin-2 in human keratinocytes in a protein kinase C-dependent manner. *Arch Dermatol Res* (2004) 295:474–81. doi:10.1007/s00403-003-0445-0
  75. Watanabe S, Kano R, Sato H, Nakamura Y, Hasegawa A. The effects of *Malassezia* yeasts on cytokine production by human keratinocytes. *J Invest Dermatol* (2001) 116:769–73. doi:10.1046/j.1523-1747.2001.01321.x
  76. Ribbing C, Engblom C, Lappalainen J, Lindstedt K, Kovanen PT, Karlsson MA, et al. Mast cells generated from patients with atopic eczema have enhanced



- levels of granule mediators and an impaired Dectin-1 expression. *Allergy* (2011) 66:110–9. doi:10.1111/j.1398-9995.2010.02437.x
77. Baroni A, Paoletti I, Ruocco E, Agozzino M, Tufano MA, Donnarumma G. Possible role of *Malassezia furfur* in psoriasis: modulation of TGF- $\beta$ 1, integrin, and HSP70 expression in human keratinocytes and in the skin of psoriasis-affected patients. *J Cutan Pathol* (2004) 31:35–42. doi:10.1046/j.0303-6987.2004.0135.x
  78. Jarvikallio A, Naukkarinen A, Harvima IT, Aalto ML, Horsmanheimo M. Quantitative analysis of tryptase- and chymase-containing mast cells in atopic dermatitis and nummular eczema. *Br J Dermatol* (1997) 136:871–7. doi:10.1111/j.1365-2133.1997.tb03927.x
  79. Selander C, Engblom C, Nilsson G, Scheynius A, Andersson CL. TLR2/MyD88-dependent and -independent activation of mast cell IgE responses by the skin commensal yeast *Malassezia sympodialis*. *J Immunol* (2009) 182:4208–16. doi:10.4049/jimmunol.0800885
  80. Gehrmann U, Qazi KR, Johansson C, Hultenby K, Karlsson M, Lundeberg L, et al. Nanovesicles from *Malassezia sympodialis* and host exosomes induce cytokine responses – novel mechanisms for host-microbe interactions in atopic eczema. *PLoS One* (2011) 6:e21480. doi:10.1371/journal.pone.0021480
  81. Ashbee HR, Bond R. *Malassezia* species and immunity: host–pathogen interactions. In: Boekhout T, Mayser P, Guého-Kellermann E, Velegraki A, editors. *Malassezia and the Skin: Science and Clinical Practice*. Berlin, Heidelberg: Springer (2010). p. 139–73.
  82. Valli JL, Williamson A, Sharif S, Rice J, Shewen PE. In vitro cytokine responses of peripheral blood mononuclear cells from healthy dogs to distemper virus, *Malassezia* and Toxocara. *Vet Immunol Immunopathol* (2010) 134:218–29. doi:10.1016/j.vetimm.2009.09.023
  83. Kang MH, Kim HJ, Jang HJ, Park HM. Sensitization rates of causative allergens for dogs with atopic dermatitis: detection of canine allergen-specific IgE. *J Vet Sci* (2014) 15:545–50. doi:10.4142/jvs.2014.15.4.545
  84. Chen TA, Halliwell RE, Pemberton AD, Hill PB. Identification of major allergens of *Malassezia pachydermatis* in dogs with atopic dermatitis and *Malassezia* overgrowth. *Vet Dermatol* (2002) 13:141–50. doi:10.1046/j.1365-3164.2002.00291.x
  85. Johansson C, Sandstrom MH, Bartosik J, Sarnhult T, Christiansen J, Zargari A, et al. Atopy patch test reactions to *Malassezia* allergens differentiate subgroups of atopic dermatitis patients. *Br J Dermatol* (2003) 148:479–88. doi:10.1046/j.1365-2133.2003.05093.x
  86. Glatz M, Buchner M, von Bartenwerffer W, Schmid-Grendelmeier P, Worm M, Hedderich J, et al. *Malassezia* spp.-specific immunoglobulin E level is a marker for severity of atopic dermatitis in adults. *Acta Derm Venereol* (2015) 95:191–6. doi:10.2340/00015555-1864
  87. Zhang E, Tanaka T, Tajima M, Tsuboi R, Kato H, Nishikawa A, et al. Anti-*Malassezia*-specific IgE antibodies production in Japanese patients with head and neck atopic dermatitis: relationship between the level of specific ige antibody and the colonization frequency of cutaneous *Malassezia* species and clinical severity. *J Allergy (Cairo)* (2011) 2011:645670. doi:10.1155/2011/645670
  88. Johansson C, Ahlberg N, Andersson A, Lundeberg L, Karlsson MA, Scheynius A, et al. Elevated peripheral allergen-specific T cell response is crucial for a positive atopy patch test reaction. *Int Arch Allergy Immunol* (2009) 150:51–8. doi:10.1159/000210380
  89. Levy JMS, Magro C. Atrophying pityriasis versicolor as an idiosyncratic T cell-mediated response to *Malassezia*: a case series. *J Am Acad Dermatol* (2017) 76:730–5. doi:10.1016/j.jaad.2016.08.062
  90. Eyerich S, Eyerich K, Pennino D, Carbone T, Nasorri F, Pallotta S, et al. Th22 cells represent a distinct human T cell subset involved in epidermal immunity and remodeling. *J Clin Invest* (2009) 119:3573–85. doi:10.1172/JCI40202
  91. Cavani A, Pennino D, Eyerich K. Th17 and Th22 in skin allergy. *Chem Immunol Allergy* (2012) 96:39–44. doi:10.1159/000331870
  92. Di Cesare A, Di Meglio P, Nestle FO. The IL-23/Th17 axis in the immunopathogenesis of psoriasis. *J Invest Dermatol* (2009) 129:1339–50. doi:10.1038/jid.2009.59
  93. Balaji H, Heratizadeh A, Wichmann K, Niebuhr M, Cramer R, Scheynius A, et al. *Malassezia sympodialis* thioredoxin-specific T cells are highly cross-reactive to human thioredoxin in atopic dermatitis. *J Allergy Clin Immunol* (2011) 128: 92–9.e4. doi:10.1016/j.jaci.2011.02.043
  94. Esaki H, Brunner PM, Renert-Yuval Y, Czarnowicki T, Huynh T, Tran G, et al. Early-onset pediatric atopic dermatitis is TH2 but also TH17 polarized in skin. *J Allergy Clin Immunol* (2016) 138:1639–51. doi:10.1016/j.jaci.2016.07.013
  95. Sparber F, LeibundGut-Landmann S. Interleukin 17-mediated host defense against *Candida albicans*. *Pathogens* (2015) 4:606–19. doi:10.3390/pathogens4030606
  96. Gaitanis G, Magiatis P, Hantschke M, Bassukas ID, Velegraki A. The *Malassezia* genus in skin and systemic diseases. *Clin Microbiol Rev* (2012) 25:106–41. doi:10.1128/CMR.00021-11
  97. Selander C, Zargari A, Mollby R, Rasool O, Scheynius A. Higher pH level, corresponding to that on the skin of patients with atopic eczema, stimulates the release of *Malassezia sympodialis* allergens. *Allergy* (2006) 61:1002–8. doi:10.1111/j.1398-9995.2006.01108.x
  98. Schmid-Grendelmeier P, Fluckiger S, Disch R, Trautmann A, Wuthrich B, Blaser K, et al. IgE-mediated and T cell-mediated autoimmunity against manganese superoxide dismutase in atopic dermatitis. *J Allergy Clin Immunol* (2005) 115:1068–75. doi:10.1016/j.jaci.2005.01.065
  99. Glaser AG, Limacher A, Fluckiger S, Scheynius A, Scapozza L, Cramer R. Analysis of the cross-reactivity and of the 1.5 A crystal structure of the *Malassezia sympodialis* Mala s 6 allergen, a member of the cyclophilin pan-allergen family. *Biochem J* (2006) 396:41–9. doi:10.1042/BJ20051708
  100. Youn SW, Na JI, Choi SY, Huh CH, Park KC. Regional and seasonal variations in facial sebum secretions: a proposal for the definition of combination skin type. *Skin Res Technol* (2005) 11:189–95. doi:10.1111/j.1600-0846.2005.00119.x
  101. Kim SY, Kim SH, Kim SN, Kim AR, Kim YR, Kim MJ, et al. Isolation and identification of *Malassezia* species from Chinese and Korean patients with seborrheic dermatitis and in vitro studies on their bioactivity on sebaceous lipids and IL-8 production. *Mycoses* (2016) 59:274–80. doi:10.1111/myc.12456
  102. Schonherr FA, Sparber F, Kirchner FR, Guiducci E, Trautwein-Weidner K, Gladiator A, et al. The intraspecies diversity of *C. albicans* triggers qualitatively and temporally distinct host responses that determine the balance between commensalism and pathogenicity. *Mucosal Immunol* (2017) 10:1335–50. doi:10.1038/mi.2017.2
  103. Zhu Y, Engstrom PG, Tellgren-Roth C, Baudo CD, Kennell JC, Sun S, et al. Proteogenomics produces comprehensive and highly accurate protein-coding gene annotation in a complete genome assembly of *Malassezia sympodialis*. *Nucleic Acids Res* (2017) 45:2629–43.

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Sparber and LeibundGut-Landmann. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.